

# Advanced Plant Health and Broad Spectrum Insect Control for Soybeans

## Benefits of Priaxor™ Fungicide


- Advanced Plant Health benefits
- Longest lasting protection
- Additional mode of action for resistance management
- More consistent performance for maximum yield potential

Increase your soybean yield with a combination of Priaxor fungicide plus Fastac insecticide

## Benefits of Fastac™ Insecticide

- Broad spectrum insect control
- Labelled on diverse crops
- Globally marketed

## 2010–2012 Replicated Trial Summary



Summary of replicated Tech Service and Biology data. Priaxor 4 fl oz/A, Fastac Insecticide 3.2 fl oz/A, R3 Applications.

## Priaxor™ Best Use Recommendations

- Use Rate: 4 fl oz/A
- Ground and aerial applications
- Priaxor may be applied at all vegetative and reproductive stages of soybean development
- The optimal application timing is from R2 stage through R4
- Priaxor may be tank mixed with other products, such as insecticides
- Adjuvant Flexible

### Target Diseases

- |                        |                                    |
|------------------------|------------------------------------|
| ■ Alternaria leaf spot | ■ Frogeye leaf spot                |
| ■ Anthracnose          | ■ Pod and stem blight              |
| ■ Asian soybean rust   | ■ Rhizoctonia aerial web blight    |
| ■ Brown spot           | ■ Sclerotinia blight (suppression) |
| ■ Cercospora blight    |                                    |

## Fastac™ Best Use Recommendations

- Use Rate: 2.8–3.8 fl/oz A
  - Maximum Cumulative Rate/Season: 11.4 fl/oz A
  - PHI: 21 Days
  - REI: 12 Hours
- Fastac is a restricted use pesticide

### Target Soybean Pests

- |                          |                                 |
|--------------------------|---------------------------------|
| ■ Alfalfa caterpillar    | ■ Mexican bean beetle           |
| ■ Armyworm               | ■ Plant bug spp.                |
| ■ Bean leaf beetle       | ■ Potato leafhopper             |
| ■ Colorado potato beetle | ■ Soybean aphid                 |
| ■ Corn earworm           | ■ Three-cornered alfalfa hopper |
| ■ Cucumber beetle        | ■ Velvetbean caterpillar        |
| ■ European corn borer    | ■ Looper spp.                   |
| ■ Green cloverworm       | ■ Stink bug spp.                |
| ■ Imported cabbageworm   |                                 |
| ■ Leaf skeletonizer spp. |                                 |

Ask your retailer about Priaxor and Fastac today.

Optimal application is R2–R4 growth stage.

**Priaxor™ + Fastac™ EC**  
Xemium® Brand Fungicide    Insecticide

**Priaxor™**  
Xemium® Brand Fungicide

**Fastac™ EC**  
Insecticide

Always read and follow label directions.  
 Priaxor and Fastac are trademarks of BASF.  
 ©2013 BASF Corporation. All Rights Reserved. APN #XX-XX-XXX

**BASF**  
The Chemical Company