



## Gramoxone<sup>®</sup> SL 2.0

syngenta<sup>®</sup>

### Get Ahead of Weeds Early

**Gramoxone<sup>®</sup> SL 2.0** herbicide delivers excellent early weed control for resistance-conscious growers. A tool that works even in cool, wet weather and is compatible with various water qualities and calcium-containing nitrogen fertilizers including CAN17, Gramoxone SL 2.0 brings convenience, flexibility and dependability for growers who want to get a head start on weed control.

#### Reliable Tool to Help Fight Herbicide Resistance

As a Resistance Fighter™ brand Gramoxone SL 2.0 brings another important mode of action to growers' weed management arsenal. When combined with residual herbicides like Reflex<sup>®</sup>, Prefix<sup>®</sup>, Sequence<sup>®</sup>, Boundary<sup>®</sup> or Dual Magnum<sup>®</sup>, Gramoxone SL 2.0 adds outstanding contact, broad-spectrum control of even the most difficult-to-control weeds. It provides fast-acting results in as little as 48 hours and is rainfast upon drying. A versatile tool, Gramoxone SL 2.0 can be used at burndown, preplant and/or pre-emergence to provide quality weed control while preserving the longevity of existing weed management tools.

#### Product Attributes

- Broad-spectrum control and proven efficacy on difficult-to-control weeds
- Fast-acting results in as little as 48 hours
- Excellent rotational partner to help preserve glyphosate efficacy and reduce the likelihood of resistance
- Provides effective control even in adverse weather conditions
- Rainfast upon drying for reliable control
- Protects crop integrity by not causing damage through root uptake
- Tank mix flexibility adds versatility to weed management programs



### Use Recommendations for Cotton

- For burndown use, apply to emerged weeds between one and six inches tall.
- Apply preplant or pre-emergence application at 2.5 to 4 pt/A, depending on weed height.
- Apply post-emergence at 32 fl oz/A with hooded sprayer.
- For necessary coverage of foliage at burndown, apply the mixture at a minimum volume of 10 gallons/A with flat-fan nozzles at 30 to 50 psi (AI nozzles not advised).
- Always add non-ionic surfactant (NIS) at 0.25% v/v or crop oil concentrate (COC) at 1% v/v.

### Use Recommendations for Soybean

- Apply at a rate of 2-4 pt/A, dependent on weed height at application.
- Apply to emerged weeds between 1-6 inches tall.
- Use as a broadcast spray before, during or after planting, but before crop emergence.
- Use a minimum of 10 gals/A.
- Use a non-ionic surfactant or crop oil concentrate.

### Use Recommendations for Peanut

- Apply 8 to 16 fl oz/A depending on weed target and height.
- Gramoxone SL 2.0 should be applied to emerged weeds when they are small (1 to 6 inches).
- Apply to control or suppress emerged annual grass and broadleaf weeds at ground crack.
- A second application may be made up to 28 days after ground crack.
- Do not apply by air.

## Target Weeds Controlled \*

Barnyardgrass

Florida beggarweed

Italian ryegrass

Johnsongrass

Palmer amaranth

Morningglory

Sicklepod

\*Refer to product label for complete list of weeds controlled.



©2012 Syngenta. Important: Always read and follow all bag tag and label instructions before buying or using Syngenta products. The instructions contain important conditions of sale, including limitations of warranty and remedy. Some crop protection products and seed treatments may not be registered for sale or use in all states or counties. Please check with your state or local extension service before buying or using Syngenta products. Gramoxone SL 2.0 is

a Restricted Use Pesticide. Boundary®, Dual Magnum®, Gramoxone®, Prefix®, Reflex®, Resistance Fighter™, Sequence®, the Alliance Frame, the Purpose icon and the Syngenta logo are trademarks of a Syngenta Group Company.